

Gauges and functional measures in quantum gravity II: Higher derivative gravity

N. Ohta,^{a,*} R. Percacci^{b,c,†} and A. D. Pereira^{d,‡}

^a*Department of Physics, Kindai University, Higashi-Osaka, Osaka 577-8502, Japan*

^b*International School for Advanced Studies, via Bonomea 265, 34136 Trieste, Italy*

^c*INFN, Sezione di Trieste, Italy*

^d*UERJ – Universidade do Estado do Rio de Janeiro, Departamento de Física Teórica, Rua São Francisco Xavier 524, 20550-013, Maracanã, Rio de Janeiro, Brasil*

Abstract

We compute the one-loop divergences in a higher-derivative theory of gravity including Ricci tensor squared and Ricci scalar squared terms, in addition to the Hilbert and cosmological terms, on an (generally off-shell) Einstein background. We work with a two-parameter family of path integral measures, corresponding to different ways of parametrizing the graviton field, and a two-parameter family of gauges. Only the logarithmic divergences are independent of these choices, but there are some choices of gauge or parametrization that reduce the dependence on the remaining parameters. The results are invariant under a recently discovered “duality” that involves the replacement of the densitized metric by a densitized inverse metric as the fundamental quantum variable.

*e-mail address: ohtan@phys.kindai.ac.jp

†e-mail address: percacci@sissa.it

‡e-mail address: duarte763@gmail.com

1 Introduction

In a previous paper [1], hereafter referred to as I, we have examined the properties of quantum General Relativity (GR - the theory containing only terms up to second derivatives in the action) in a general four-parameter family of gauges and parametrizations. In this paper, we would like to extend the analysis to Higher-Derivative Gravity (HDG).

By HDG, we will always mean the theory of gravity based on the metric as fundamental variable and an action that contains up to four derivatives.¹ This theory is important because in four dimensions it is power-counting renormalizable [5] and asymptotically free [6, 7, 8]. In spite of these appealing properties, it has never been accepted as a viable fundamental theory for gravity, because of the presence of ghosts in perturbation theory. Over time, there have been many proposals trying to circumvent this problem. Among these let us mention here the following possibilities:

- The mass of the ghost is not a fixed parameter but is rather subject to strong (quadratic) running above the Planck threshold. Then, the equation for the pole mass $m_{phys}^2 = m^2(k = m_{phys})$ (where $m(k)$ is the running mass) may not have a solution [6, 9, 10].
- The ghost may be an artifact of expanding around the wrong vacuum. The true vacuum of quadratic gravity (in the presence also of a Hilbert term) is not flat space but rather a kind of wave with wavelength of the order of the Planck length [11].
- The quadratic term is one of an infinite series and the sum of the series is a function that has no massive ghost pole. The ghost pole is an artifact of Taylor expanding this function to second order (see the aforementioned papers on non-local gravity and also [12]).

For further proposals see also [13, 14, 15]. So far, none of these arguments has convinced the community at large, so the issue of the ghosts remains open for the time being. However, the hope that a way out may exist has generated new interest in these theories in recent times, also among particle physicists [16, 17, 18]. It seems therefore appropriate to keep investigating the quantum properties of these theories.

In this paper we will extend previous results in several directions. Due to the complicated structure of the theory, calculations of one-loop divergences in HDG have usually been performed with a special four-derivative gauge-fixing term such that the four-derivative part of the Hessian is proportional to the square of the Laplacian. In this paper we will calculate the off-shell gauge-dependence of the one-loop divergences by using the more conventional second-derivative gauge-fixing term that is commonly used in quantum GR, depending on two parameters a and b , or a four-derivative variant of the same gauge fixing, containing an extra power of a Laplacian.

As another generalization, we assume that the functional measure is not defined on the metric but on a densitized metric

$$\gamma_{\mu\nu} = g_{\mu\nu} \left(\sqrt{\det g_{\mu\nu}} \right)^w, \quad (1.1)$$

or densitized inverse metric

$$\gamma^{\mu\nu} = g^{\mu\nu} \left(\sqrt{\det g^{\mu\nu}} \right)^{-w} = g^{\mu\nu} \left(\sqrt{\det g_{\mu\nu}} \right)^w, \quad (1.2)$$

¹There has been recently some progress on theories that contain more than four, and possibly infinitely many derivatives [2, 3, 4]. These also deserve the name HDG, but we shall not consider them here.

with weight w . Furthermore we allow the quantum theory to depend on another parameter ω that interpolates continuously between the linear background-field expansion (for $\omega = 0$)

$$\gamma_{\mu\nu} = \bar{\gamma}_{\mu\nu} + \hat{h}_{\mu\nu} , \quad (1.3)$$

the exponential background-field expansions (for $\omega = 1/2$)

$$\gamma_{\mu\nu} = \bar{\gamma}_{\mu\rho}(e^{\hat{h}})^{\rho}_{\nu} \quad \text{or} \quad \gamma^{\mu\nu} = (e^{-\hat{h}})^{\mu}_{\rho}\bar{\gamma}^{\rho\nu} . \quad (1.4)$$

and the linear expansion around the inverse densitized metric (for $\omega = 1$)

$$\gamma^{\mu\nu} = \bar{\gamma}^{\mu\nu} - \hat{h}^{\mu\nu} . \quad (1.5)$$

The properties of quantum GR in this general four-parameter family of gauges and parametrizations have been investigated in an accompanying paper [1].

Here we extend the analysis of I to HDG. We now have four independent couplings instead of two, so the expressions for the divergences are in general much more complicated than in I and so is the interpretation of the results. The expressions simplify somewhat, and are reported explicitly, in two limits: the “four derivative gravity (4DG) limit” in which the Einstein-Hilbert terms can be neglected relative to the curvature squared terms, and the “Einstein-Hilbert (EH) limit” where the opposite holds. In the EH limit, the action is the same as the one considered in I, but the analysis that we perform here differs in two respects: first, we choose a more general Einstein background, instead of the maximally symmetric background of I. This allows us to discriminate two divergent terms quadratic in curvature, rather than a single one as in I. Secondly, the cutoff in each spin sector is chosen to depend on the corresponding Lichnerowicz Laplacian, rather than the Bochner Laplacian $-\nabla^2$ as in I. Thus, comparison with I yields some information on the cutoff-dependence of these non-universal results. Still, we find that in this limit the qualitative picture is the same.

In four dimensions and in the 4DG-limit, all divergences are universal, i.e. independent of both gauge and parametrization. In particular, the logarithmic divergences are related to the well-known universal beta functions of HDG [8].

As in I, we find that all divergences are invariant under a “duality” transformation that consists in the replacement

$$\begin{aligned} \omega &\rightarrow 1 - \omega, \\ w &\rightarrow w + \frac{4}{d} . \end{aligned} \quad (1.6)$$

It was shown in I that this transformation leaves the functional measure invariant.

We conclude this introduction by listing the contents of the following sections. Section 2 contains details of our calculations of one-loop divergences. In section 3 we give a formal proof (at the level of determinants) that the effective action of HDG in the 4DG-limit is gauge-independent. The main results are presented in section 4. In section 5, we discuss how the results can be obtained for $d = 4$ conformal gravity. In section 6, we point out that our results show that the duality found in our previous paper I is valid in HDG as well. Section 5 contains a discussion and conclusions. In the appendix, we summarize the heat kernel coefficients for Lichnerowicz Laplacians on an Einstein manifold.

2 The one-loop effective action

2.1 The HDG actions and their equations of motion

In this paper we will consider actions of the general form

$$S = \int d^d x \sqrt{\mp g} \left[\pm Z_N (R - 2\Lambda) + \alpha R^2 + \beta R_{\mu\nu}^2 \right], \quad (2.1)$$

where $Z_N = 1/(16\pi G)$, Λ is the cosmological constant, α, β are the higher derivative couplings. The upper sign refers to Minkowski signature, the lower one to Euclidean signature. This is not the most general HDG action, because we omit a term $\gamma R_{\mu\nu\rho\sigma}^2$, however in $d \leq 4$ this is not a very strong restriction, because the ‘‘Gauss-Bonnet’’ or ‘‘Euler’’ combination

$$G = R_{\mu\nu\alpha\beta}^2 - 4R_{\mu\nu}^2 + R^2, \quad (2.2)$$

is either zero (in $d = 2, 3$) or a total derivative (in $d = 4$). Using the identity for the Weyl tensor $C_{\mu\nu\rho\sigma}$:

$$C^2 \equiv C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = R_{\mu\nu\alpha\beta}^2 - \frac{4}{d-2} R_{\mu\nu}^2 + \frac{2}{(d-1)(d-2)} R^2, \quad (2.3)$$

one has

$$C^2 = G + \frac{2(d-3)}{d-2} W; \quad W = 2R_{\mu\nu}^2 - \frac{d}{2(d-1)} R^2. \quad (2.4)$$

Thus, modulo terms proportional to G , we can replace $\alpha R^2 + \beta R_{\mu\nu}^2$ by

$$\frac{1}{2\lambda} W + \frac{1}{\xi} R^2. \quad (2.5)$$

The integral of W , like the integral of C^2 , is Weyl-invariant in $d = 4$.

The equations of motion are

$$\pm Z_N (G_{\mu\nu} + \Lambda g_{\mu\nu}) + \alpha E_{\mu\nu}^{(1)} + \beta E_{\mu\nu}^{(2)} = 0, \quad (2.6)$$

where $G_{\mu\nu}$ is the Einstein tensor and

$$\begin{aligned} E_{\mu\nu}^{(1)} &= 2R R_{\mu\nu} - 2\nabla_\mu \nabla_\nu R + g_{\mu\nu} \left(2\Box R - \frac{1}{2} R^2 \right), \\ E_{\mu\nu}^{(2)} &= 2R_{\mu\lambda} R_\nu^\lambda - 2\nabla^\lambda \nabla_{(\mu} R_{\nu)\lambda} + \Box R_{\mu\nu} + \frac{1}{2} (\Box R - R_{\lambda\rho}^2) g_{\mu\nu}, \end{aligned} \quad (2.7)$$

Let us observe that in the case $Z_N = 0$ and $d = 4$ the Einstein condition

$$R_{\mu\nu} = \frac{1}{d} R g_{\mu\nu}, \quad (2.8)$$

is enough to fulfil the equations of motion. The scalar curvature remains an undetermined constant, since the action is scale-invariant. On the other hand when $Z_N \neq 0$ the equations of motion together with the Einstein condition further require

$$\mathcal{P} \equiv \frac{4-d}{2} \left(\alpha + \frac{\beta}{d} \right) R^2 \mp \frac{d-2}{2} Z_N \left(R - \frac{2d}{d-2} \Lambda \right) = 0. \quad (2.9)$$

In particular, in four dimensions the higher-derivative terms do not contribute and the equation of motion is the same as the trace of the Einstein equations with cosmological constant:

$$E \equiv R - \frac{2d}{d-2} \Lambda = 0. \quad (2.10)$$

2.2 Quadratic expansion

The action has to be thought of as a functional of the quantum field $\gamma_{\mu\nu}$ or $\gamma^{\mu\nu}$ defined as in (1.1) or (1.2). For $w \neq -1/d$ these relations can be inverted to yield

$$g_{\mu\nu} = \gamma_{\mu\nu} (\det(\gamma_{\mu\nu}))^m ; \quad g^{\mu\nu} = \gamma^{\mu\nu} (\det(\gamma_{\mu\nu}))^{-m} , \quad (2.11)$$

where

$$\frac{w}{2} = -\frac{m}{1+dm} \quad \text{or} \quad \frac{w}{2} = \frac{m}{1+dm} , \quad (2.12)$$

respectively. Conversely, $m = -\frac{w/2}{1+dw/2}$ for (1.1) and $m = \frac{w/2}{1-dw/2}$ for (1.2). We observe that the relation between m and $w/2$ is an involution. We choose to treat m as an independent free parameter. All dependence on m can be translated into a dependence on w if needed, using the preceding formulas.

The quantum field is expanded as in (1.3) or (1.4) or (1.5), and for the calculation of the one-loop divergences we need the terms to second order in the fluctuation. As explained in [1], we can start from the quadratic expansion of the action in $\delta g_{\mu\nu}$, which has been given in detail in [19], and then use

$$\delta g_{\mu\nu} = \delta g_{\mu\nu}^{(1)} + \delta g_{\mu\nu}^{(2)} + \dots , \quad (2.13)$$

where $\delta g_{\mu\nu}^{(n)}$ contains n powers of the tensor fluctuation

$$h_{\mu\nu} = (\det \bar{\gamma})^m \hat{h}_{\mu\nu} . \quad (2.14)$$

For all four types of expansion considered here, we have

$$\begin{aligned} \delta g_{\mu\nu}^{(1)} &= h_{\mu\nu} + m \bar{g}_{\mu\nu} h , \\ \delta g_{\mu\nu}^{(2)} &= \omega h_{\mu\rho} h^\rho{}_\nu + m h h_{\mu\nu} + m \left(\omega - \frac{1}{2} \right) \bar{g}_{\mu\nu} h^{\alpha\beta} h_{\alpha\beta} + \frac{1}{2} m^2 \bar{g}_{\mu\nu} h^2 . \end{aligned} \quad (2.15)$$

The choice $\omega = 0$ corresponds to the linear expansion of metric (1.3), $\omega = 1/2$ corresponds to the exponential expansion (1.4) (and it does not matter if one starts from the metric or from the inverse metric) and $\omega = 1$ corresponds to the linear expansion of the inverse metric (1.5).

2.3 Lichnerowicz Laplacians

The evaluation of the one-loop divergences is based on the knowledge of the coefficients of the small-time expansion of the heat kernel of the kinetic operator appearing in the gauge-fixed Hessian. We will denote $\bar{\nabla}$ the covariant derivative defined by the background metric $\bar{g}$, $\square = \bar{\nabla}^2$ the d'Alembertian and $-\bar{\nabla}^2$ its Euclidean analog, known as Bochner Laplacian. In GR, the expansion of the action generates non-minimal terms that can be eliminated by choosing the de Donder gauge. The rest can be written in terms of Laplacians. Similarly, the second variation of the HDG action contains non-Laplacian terms such as $\bar{\nabla}_\mu \bar{\nabla}_\nu \bar{\nabla}_\rho \bar{\nabla}_\sigma$, $\square \bar{\nabla}_\mu \bar{\nabla}_\nu \bar{g}_{\rho\sigma} + \bar{g}_{\mu\nu} \bar{\nabla}_\rho \bar{\nabla}_\sigma \square$ and $\bar{g}_{\nu\sigma} \bar{\nabla}_\mu \square \bar{\nabla}_\rho + \bar{g}_{\mu\rho} \bar{\nabla}_\nu \square \bar{\nabla}_\sigma$, acting on $h_{\rho\sigma}$. The heat kernel expansion for such operators is not known, so we will use the York decomposition to rewrite the Hessian as minimal operators acting on fields with definite spin. For this, it is essential to rewrite all occurrences of $\square$ in terms of the Lichnerowicz Laplacians acting on spin-zero, spin-one and spin-two fields, which are defined (in Euclidean signature) as follows:

$$\begin{aligned} \Delta_{L0} \phi &= -\bar{\nabla}^2 \phi , \\ \Delta_{L1} A_\mu &= -\bar{\nabla}^2 A_\mu + \bar{R}_\mu{}^\rho A_\rho , \\ \Delta_{L2} h_{\mu\nu} &= -\bar{\nabla}^2 h_{\mu\nu} + \bar{R}_\mu{}^\rho h_{\rho\nu} + \bar{R}_\nu{}^\rho h_{\mu\rho} - \bar{R}_{\mu\rho\nu\sigma} h^{\rho\sigma} - \bar{R}_{\mu\rho\nu\sigma} h^{\sigma\rho} . \end{aligned} \quad (2.16)$$

These operators have the following useful properties:

$$\Delta_{L1}\bar{\nabla}_\mu\phi = \bar{\nabla}_\mu\Delta_{L0}\phi, \quad (2.17)$$

$$\bar{\nabla}_\mu\Delta_{L1}\xi^\mu = \Delta_{L0}\bar{\nabla}_\mu\xi^\mu, \quad (2.18)$$

$$\Delta_{L2}(\bar{\nabla}_\mu\bar{\nabla}_\nu\phi) = \bar{\nabla}_\mu\bar{\nabla}_\nu\Delta_{L0}\phi, \quad (2.19)$$

$$\Delta_{L2}(\bar{\nabla}_\mu\xi_\nu + \bar{\nabla}_\nu\xi_\mu) = \bar{\nabla}_\mu\Delta_{L1}\xi_\nu + \bar{\nabla}_\nu\Delta_{L1}\xi_\mu, \quad (2.20)$$

$$\Delta_{L2}g_{\mu\nu}\phi = g_{\mu\nu}\Delta_{L0}\phi. \quad (2.21)$$

The York decomposition leads to significant simplifications when the background metric is an Einstein space, i.e. satisfies the condition (2.8). (As discussed in section 2.1, this is not enough to put the background on-shell.) In the following we will always assume that the background is Einstein, but not that it satisfies (2.10).

Now consider the second variations of the curvature squared terms in the action, as given in eqs. (B.6) and (B.7) in [19]. The terms containing the Riemann tensor can be combined with certain terms containing $\square$ to give Lichnerowicz Laplacians. Further using (2.8), one arrives at

$$\begin{aligned} & \alpha\delta g^{\mu\nu} \left[\bar{\nabla}_\mu\bar{\nabla}_\nu\bar{\nabla}_\alpha\bar{\nabla}_\beta - 2\bar{g}_{\mu\nu}\square\bar{\nabla}_\alpha\bar{\nabla}_\beta + \bar{g}_{\mu\nu}\bar{g}_{\alpha\beta}\square^2 \right. \\ & - \bar{g}_{\nu\beta}\bar{R}\bar{\nabla}_\mu\bar{\nabla}_\alpha + \frac{d-2}{2}\bar{R}\bar{g}_{\mu\nu}\bar{\nabla}_\alpha\bar{\nabla}_\beta + \frac{4-d}{2d}\bar{g}_{\mu\nu}\bar{g}_{\alpha\beta}\bar{R}\square - \frac{1}{2}\bar{g}_{\mu\alpha}\bar{g}_{\nu\beta}\bar{R}\Delta_{L2} \\ & \left. + \left(\frac{1}{d^2} - \frac{1}{d} + \frac{1}{8}\right)\bar{R}^2\bar{g}_{\mu\nu}\bar{g}_{\alpha\beta} + \left(\frac{2}{d} - \frac{1}{4}\right)\bar{R}^2\bar{g}_{\mu\alpha}\bar{g}_{\nu\beta} \right] \delta g^{\alpha\beta}, \end{aligned} \quad (2.22)$$

and

$$\begin{aligned} & \beta\delta g^{\mu\nu} \left[\frac{1}{2}\bar{\nabla}_\mu\bar{\nabla}_\nu\bar{\nabla}_\alpha\bar{\nabla}_\beta - \frac{1}{2}\bar{g}_{\mu\nu}\square\bar{\nabla}_\alpha\bar{\nabla}_\beta - \frac{1}{2}\bar{g}_{\nu\beta}\bar{\nabla}_\mu\square\bar{\nabla}_\alpha + \frac{1}{4}\bar{g}_{\mu\nu}\bar{g}_{\alpha\beta}\square^2 \right. \\ & + \frac{1}{4}\bar{g}_{\mu\alpha}\bar{g}_{\nu\beta}\Delta_{L2} \left(\Delta_{L2} - \frac{6}{d}\bar{R} \right) - \frac{3}{2d}\bar{R}\bar{g}_{\nu\beta}\bar{\nabla}_\mu\bar{\nabla}_\alpha + \frac{1}{d}\bar{R}\bar{g}_{\mu\nu}\bar{\nabla}_\alpha\bar{\nabla}_\beta - \frac{1}{4d}\bar{R}\bar{g}_{\mu\nu}\bar{g}_{\alpha\beta}\square \\ & \left. + \frac{12-d}{4d^2}\bar{R}^2\bar{g}_{\mu\nu}\bar{g}_{\alpha\beta} + \frac{d-8}{8d^2}\bar{R}^2\bar{g}_{\mu\alpha}\bar{g}_{\nu\beta} \right] \delta g^{\alpha\beta}. \end{aligned} \quad (2.23)$$

All tensor structures are provided by the background metric. We note that this procedure of eliminating the Riemann tensor does not work in the case of the second variation of the Riemann squared term, which is the reason why we do not consider such a term.

2.4 York decomposition

The York decomposition is defined by

$$h_{\mu\nu} = h_{\mu\nu}^{TT} + \bar{\nabla}_\mu\xi_\nu + \bar{\nabla}_\nu\xi_\mu + \bar{\nabla}_\mu\bar{\nabla}_\nu\sigma - \frac{1}{d}\bar{g}_{\mu\nu}\bar{\nabla}^2\sigma + \frac{1}{d}\bar{g}_{\mu\nu}h, \quad (2.24)$$

where $h_{\mu\nu}^{TT}$ is transverse and tracefree, and $\hat{\xi}_\mu$ is transverse. We will use

$$\hat{\xi}_\mu = \sqrt{-\bar{\nabla}^2 - \frac{\bar{R}}{d}}\xi_\mu \quad ; \quad \hat{\sigma} = \sqrt{-\bar{\nabla}^2}\sqrt{-\bar{\nabla}^2 - \frac{\bar{R}}{d-1}}\sigma. \quad (2.25)$$

Employing (2.24) and (2.25) and assuming that the background is an Einstein space, one finds

$$\int d^d x \sqrt{\bar{g}} h_{\mu\nu} h^{\mu\nu} = \int d^d x \sqrt{\bar{g}} \left[h_{\mu\nu}^{TT} h^{TT\mu\nu} + 2\hat{\xi}_\mu\hat{\xi}^\mu + \frac{d-1}{d}\hat{\sigma}^2 + \frac{1}{d}h^2 \right]. \quad (2.26)$$

Using the properties (2.17) – (2.21), we have

$$\int d^d x \sqrt{\bar{g}} h_{\mu\nu} \Delta_{L2} h^{\mu\nu} = \int d^d x \sqrt{\bar{g}} \left[h_{\mu\nu}^{TT} \Delta_{L2} h^{TT\mu\nu} + 2\hat{\xi}_\mu \Delta_{L1} \hat{\xi}^\mu + \frac{d-1}{d} \hat{\sigma} \Delta_{L0} \hat{\sigma} + \frac{1}{d} h \Delta_{L0} h \right], \quad (2.27)$$

$$\begin{aligned} \int d^d x \sqrt{\bar{g}} h_{\mu\nu} (\Delta_{L2})^2 h^{\mu\nu} &= \int d^d x \sqrt{\bar{g}} \left[h_{\mu\nu}^{TT} (\Delta_{L2})^2 h^{TT\mu\nu} + 2\hat{\xi}_\mu (\Delta_{L1})^2 \hat{\xi}^\mu \right. \\ &\quad \left. + \frac{d-1}{d} \hat{\sigma} (\Delta_{L0})^2 \hat{\sigma} + \frac{1}{d} h (\Delta_{L0})^2 h \right], \end{aligned} \quad (2.28)$$

and so on.

Furthermore using

$$\bar{\nabla}_\alpha h^\alpha{}_\nu = \left(\bar{\nabla}^2 + \frac{\bar{R}}{d} \right) \xi_\nu + \frac{d-1}{d} \bar{\nabla}_\nu \left(\bar{\nabla}^2 + \frac{\bar{R}}{d-1} \right) \sigma + \frac{1}{d} \bar{\nabla}_\nu h, \quad (2.29)$$

and

$$\bar{\nabla}_\alpha \bar{\nabla}_\beta h^{\alpha\beta} = \frac{d-1}{d} \bar{\nabla}^2 \left(\bar{\nabla}^2 + \frac{\bar{R}}{d-1} \right) \sigma + \frac{1}{d} \bar{\nabla}^2 h, \quad (2.30)$$

we find that

$$\begin{aligned} \int d^d x \sqrt{\bar{g}} h^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\nu \bar{\nabla}_\alpha \bar{\nabla}_\beta h^{\alpha\beta} &= \int d^d x \sqrt{\bar{g}} \left[\left(\frac{d-1}{d} \right)^2 \sigma \square^2 \left(\square + \frac{\bar{R}}{d-1} \right) \sigma \right. \\ &\quad \left. + 2 \frac{d-1}{d^2} h \square^2 \left(\square + \frac{\bar{R}}{d-1} \right) \sigma + \frac{1}{d^2} h \square^2 h \right], \end{aligned} \quad (2.31)$$

$$\begin{aligned} \int d^d x \sqrt{\bar{g}} h^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\alpha h^\alpha{}_\nu &= \int d^d x \sqrt{\bar{g}} \left[-\xi_\mu \left(\square + \frac{\bar{R}}{d} \right)^2 \xi^\mu + \left(\frac{d-1}{d} \right)^2 \sigma \square \left(\square + \frac{\bar{R}}{d-1} \right)^2 \sigma \right. \\ &\quad \left. + 2 \frac{d-1}{d^2} h \square \left(\square + \frac{\bar{R}}{d-1} \right) \sigma + \frac{1}{d^2} h \square h \right], \end{aligned} \quad (2.32)$$

$$\begin{aligned} \int d^d x \sqrt{\bar{g}} h^{\mu\nu} \bar{\nabla}_\mu \square \bar{\nabla}_\alpha h^\alpha{}_\nu &= \int d^d x \sqrt{\bar{g}} \left[-\xi_\mu \square \left(\square + \frac{\bar{R}}{d} \right)^2 \xi^\mu \right. \\ &\quad + \left(\frac{d-1}{d} \right)^2 \sigma \square \left(\square + \frac{\bar{R}}{d} \right) \left(\square + \frac{\bar{R}}{d-1} \right)^2 \sigma \\ &\quad \left. + 2 \frac{d-1}{d^2} h \square \left(\square + \frac{\bar{R}}{d} \right) \left(\square + \frac{\bar{R}}{d-1} \right) \sigma + \frac{1}{d^2} h \square \left(\square + \frac{\bar{R}}{d} \right) h \right]. \end{aligned} \quad (2.33)$$

2.5 The decomposed Hessian

The expansion of the Euclidean action in powers of h has the following quadratic part

$$S^{(2)} = \int d^d x \sqrt{\bar{g}} \left[h_{\mu\nu}^{TT} H^{TT} h^{TT\mu\nu} + \hat{\xi}_\mu H^{\xi\xi} \hat{\xi}^\mu + \hat{\sigma} H^{\sigma\sigma} \hat{\sigma} + \hat{\sigma} H^{\sigma h} h + h H^{h\sigma} \hat{\sigma} + h H^{hh} h \right], \quad (2.34)$$

with

$$\begin{aligned}
H^{TT} = & \frac{1}{4}Z_N \left(\Delta_{L2} - \frac{2\bar{R}}{d} + \frac{d-2}{d}(1-2\omega)(1+dm)\bar{E} \right) \\
& + \frac{\beta}{4} \left((\Delta_{L2})^2 - \frac{6}{d}\bar{R}\Delta_{L2} + \frac{8-(d-4)(1-2\omega)(1+dm)}{d^2}\bar{R}^2 \right) \\
& - \frac{\alpha}{2}\bar{R} \left(\Delta_{L2} - \frac{4-(d-4)(1-2\omega)(1+dm)}{2d}\bar{R} \right), \tag{2.35}
\end{aligned}$$

$$H^{\xi\xi} = \frac{(1-2\omega)(1+dm)}{2d} \left[(d-2)Z_N\bar{E} - \frac{d-4}{d}(d\alpha+\beta)\bar{R}^2 \right], \tag{2.36}$$

$$\begin{aligned}
H^{\sigma\sigma} = & \frac{d-1}{d} \left\{ -\frac{d-2}{4d}Z_N(\Delta_{L0} - (1-2\omega)(1+dm)\bar{E}) \right. \\
& + \frac{d-1}{d}\alpha \left[(\Delta_{L0})^2 + \frac{d-4}{2(d-1)}\bar{R} \left(\Delta_{L0} - \frac{(1-2\omega)(1+dm)}{2}\bar{R} \right) \right] \\
& \left. + \frac{\beta}{4} \left[(\Delta_{L0})^2 + \frac{2(d-4)}{d^2}\bar{R} \left(\Delta_{L0} - \frac{(1-2\omega)(1+dm)}{2}\bar{R} \right) \right] \right\}, \tag{2.37}
\end{aligned}$$

$$\begin{aligned}
H^{\sigma h} = H^{h\sigma} = & \frac{d-1}{2d}(1+dm)\sqrt{\Delta_{L0}}\sqrt{\Delta_{L0} - \frac{\bar{R}}{d-1}} \left[-Z_N\frac{d-2}{2d} \right. \\
& \left. + \alpha\frac{2(d-1)}{d} \left(\Delta_{L0} + \frac{d-4}{2(d-1)}\bar{R} \right) + \frac{\beta}{2} \left(\Delta_{L0} + \frac{2(d-4)}{d^2}\bar{R} \right) \right], \tag{2.38}
\end{aligned}$$

$$\begin{aligned}
H^{hh} = & \frac{d-1}{d}(1+dm)^2 \left\{ -Z_N\frac{d-2}{4d} \left(\Delta_{L0} - \frac{\bar{R}}{d-1} + \frac{d-2+d^2m+4\omega}{2(d-1)(1+dm)}\bar{E} \right) \right. \\
& + \alpha\frac{d-1}{d} \left((\Delta_{L0})^2 + \frac{d-6}{2(d-1)}\bar{R} + \frac{(d-4)(d-6-4dm+d^2m+4\omega)}{8(d-1)^2(1+dm)}\bar{R}^2 \right) \\
& \left. + \frac{\beta}{4} \left((\Delta_{L0})^2 + \frac{d^2-10d+8}{d^2(d-1)}\bar{R} + \frac{(d-4)(d-6-4dm+d^2m+4\omega)}{2(d-1)d^2(1+dm)}\bar{R}^2 \right) \right\}, \tag{2.39}
\end{aligned}$$

where $\bar{E} = \bar{R} - \frac{2d\Lambda}{d-2}$.

If we define the scalar gauge-invariant degree of freedom

$$s = \frac{\sqrt{\Delta_{L0}}}{\sqrt{\Delta_{L0} - \frac{\bar{R}}{d-1}}} \hat{\sigma} + (1+dm)h, \tag{2.40}$$

the scalar sector of the Hessian can be rewritten as

$$\int d^d x \sqrt{\bar{g}} \left[s H_s^{ss} s + s H_s^{sh} h + h H_s^{hs} s + h H_s^{hh} h \right], \tag{2.41}$$

where

$$\begin{aligned}
H_s^{ss} = & \frac{d-1}{4d^3} \frac{\Delta_{L0} - \frac{\bar{R}}{d-1}}{\Delta_{L0}} \left\{ -d(d-2)Z_N \left[\Delta_{L0} - (1+dm)(1-2\omega) \left(\bar{R} - \frac{2d\Lambda}{d-2} \right) \right] \right. \\
& + 4d(d-1)\alpha \left[(\Delta_{L0})^2 + \frac{d-4}{2(d-1)}\Delta_{L0}\bar{R} - \frac{d-4}{4(d-1)}(1+dm)(1-2\omega)\bar{R}^2 \right] \\
& \left. + d^2\beta \left[(\Delta_{L0})^2 + 2\frac{d-4}{d^2}\Delta_{L0}\bar{R} - \frac{d-4}{d^2}(1+dm)(1-2\omega)\bar{R}^2 \right] \right\}, \tag{2.42}
\end{aligned}$$

$$H_s^{sh} = H_s^{hs} = -\frac{d-1}{2d^2} \frac{\Delta_{L0} - \frac{\bar{R}}{d-1}}{\Delta_{L0}} (1+dm)^2(1-2\omega)\bar{\mathcal{P}}, \tag{2.43}$$

$$\begin{aligned}
H_s^{hh} = & -\frac{1}{4d^2} \frac{(1+dm)}{\Delta_{L0}} \left[d \left(2d(d-1)m^2(2\omega-1) + dm(8\omega-3) - 4m(2\omega-1) + 4\omega-1 \right) \Delta_{L0} \right. \\
& \left. + 2(1+dm)^2(1-2\omega)\bar{R} \right] \bar{\mathcal{P}}. \tag{2.44}
\end{aligned}$$

This form of the Hessian has the virtue that all the terms that contain h are proportional to the equation of motion. This shows that the field h can be completely disregarded on shell, as one would expect of a gauge-variant variable. This however depends on the choice of basis in the space of fields and is only true when the other scalar degree of freedom is gauge-invariant.

On the other hand, this form of the Hessian has the unpleasant feature that the kinetic operator of the field s is non-local. One cannot generally absorb the non-local prefactor in a redefinition of s , because one is not allowed to perform non-local redefinitions of physical fields.

One notices, however, that the terms with the lowest power of Δ_{L0} in each of the three lines in H_s^{ss} is proportional to $(1+dm)(1-2\omega)$. Therefore, if either $\omega = 1/2$ (exponential parametrization) or $m = -1/d$ (the “unimodular” measure), each of the square brackets in (2.42) is proportional to Δ_{L0} and the Hessian of s becomes local.

We further observe that for $\omega = 1/2$ also the mixed term vanishes and the term H_s^{hh} becomes local, whereas for $m = -1/d$ all terms containing h vanish, as expected in the unimodular theory.

Furthermore, H_s^{ss} becomes local and independent of m and ω for pure four-derivative gravity ($Z_N = 0$) when $d = 4$. In this case the whole Hessian becomes just

$$\frac{1}{4}\beta h_{\mu\nu}^{TT} \left(\Delta_{L2} - \frac{\bar{R}}{2} \right) \left(\Delta_{L2} - \left(1 + \frac{2\alpha}{\beta} \right) \bar{R} \right) h^{TT\mu\nu} + \frac{3}{16}(3\alpha + \beta)s\Delta_{L0} \left(\Delta_{L0} - \frac{\bar{R}}{3} \right) s. \tag{2.45}$$

Finally we observe that in the conformal case $d = 4$, $\beta = -3\alpha$ (which corresponds to having only the term W in the action) the ss term in the Hessian vanishes too, leaving only the pure spin-two degree of freedom, with Hessian

$$\frac{1}{4}\beta h_{\mu\nu}^{TT} \left(\Delta_{L2} - \frac{\bar{R}}{2} \right) \left(\Delta_{L2} - \frac{\bar{R}}{3} \right) h^{TT\mu\nu}. \tag{2.46}$$

2.6 Two-derivative gauge fixing terms

We now turn to the discussion of the gauge-fixing and Faddeev-Popov (FP) ghost terms. In most of the following, we use the same two-derivative gauge-fixing term as in I [1]. The gauge-fixing function is defined as

$$F_\mu = \bar{\nabla}_\alpha h^\alpha{}_\mu - \frac{\bar{b}+1}{d}\bar{\nabla}_\mu h, \tag{2.47}$$

with a gauge-fixing parameter $-\infty < \bar{b} < \infty$. Here $h_{\mu\nu}$ is the tensorial quantum field defined above and the bars on the covariant derivatives means that they are calculated from the background metric $\bar{g}_{\mu\nu}$. The simplest way to derive the gauge-fixing and FP ghost terms is to use the BRST transformations which is obtained by replacing the gauge parameters with the FP ghosts. In the present case, we have

$$\delta_B g_{\mu\nu} = -\delta\lambda[g_{\rho\nu}\bar{\nabla}_\mu C^\rho + g_{\rho\mu}\bar{\nabla}_\nu C^\rho], \quad (2.48)$$

where C^μ is the FP ghost, and $\delta\lambda$ is an anticommuting parameter. The BRST transformation for other fields is derived by the requirement of the nilpotency of the transformation:

$$\delta_B C^\mu = -\delta\lambda C^\rho \bar{\nabla}_\rho C^\mu, \quad \delta_B \bar{C}_\mu = i\delta\lambda B_\mu, \quad \delta_B B_\mu = 0, \quad (2.49)$$

where $\bar{C}_\mu$ is the FP anti-ghost and B_μ is an auxiliary field which enforces the gauge-fixing condition.

In order to derive the gauge fixing and FP terms, we have to know the transformations on the fluctuation field $h_{\mu\nu}$. From the first relation in eq. (2.15), we have

$$\delta_B h_{\mu\nu} + m\bar{g}_{\mu\nu}\delta_B h + \dots = -\delta\lambda[g_{\rho\nu}\bar{\nabla}_\mu C^\rho + g_{\rho\mu}\bar{\nabla}_\nu C^\rho]. \quad (2.50)$$

Anticipating that we will compute the effective action for vanishing expectation value of $h_{\mu\nu}$, we can restrict this to the linear terms to obtain

$$\delta_B h_{\mu\nu} = -\delta\lambda \left[\bar{\nabla}_\mu C_\nu + \bar{\nabla}_\nu C_\mu - \frac{2m}{1+dm}\bar{g}_{\mu\nu}\bar{\nabla}_\rho C^\rho \right]. \quad (2.51)$$

The gauge-fixing and FP terms are then given as [22]

$$\begin{aligned} \mathcal{L}_{GF+FP}/\sqrt{\bar{g}} &= i\delta_B \left[\bar{C}_\mu \left(F^\mu + \frac{a}{2Z_{GF}} B^\mu \right) \right] / \delta\lambda \\ &= -B_\mu \left(F^\mu + \frac{a}{2Z_{GF}} B^\mu \right) + i\bar{C}_\mu \left[\bar{\nabla}_\alpha \left(\bar{\nabla}^\mu C^\alpha + \bar{\nabla}^\alpha C^\mu - \frac{2m}{1+dm}\bar{g}^{\mu\alpha}\bar{\nabla}_\rho C^\rho \right) \right. \\ &\quad \left. - \frac{1+\bar{b}}{d}\bar{\nabla}^\mu \left(\frac{2}{1+dm}\bar{\nabla}_\rho C^\rho \right) \right] \\ &= -\frac{a}{2Z_{GF}}\tilde{B}_\mu\tilde{B}^\mu + \frac{Z_{GF}}{2a}F_\mu F^\mu + i\bar{C}_\mu\Delta^{(gh)\mu}{}_\nu C^\nu, \end{aligned} \quad (2.52)$$

and we have set $\bar{b} = (1+dm)b$ and defined

$$\tilde{B}_\mu = B_\mu + \frac{Z_{GF}}{a}F_\mu, \quad \Delta_{\mu\nu}^{(gh)} \equiv \bar{g}_{\mu\nu}\square + \left(1 - 2\frac{b+1}{d}\right)\bar{\nabla}_\mu\bar{\nabla}_\nu + \bar{R}_{\mu\nu}. \quad (2.53)$$

Here a is another dimensionless gauge parameter, and Z_{GF} is a parameter with dimension $d-2$. Since the B_μ field involves no derivatives, we can simply integrate it out and then we are left with the gauge-fixing and FP ghost terms.

Using the York decomposition,

$$F_\mu = -\left(\Delta_{L1} - \frac{2\bar{R}}{d}\right)\xi_\mu - \bar{\nabla}_\mu \left(\frac{d-1}{d}\left(\Delta_{L0} - \frac{\bar{R}}{d-1}\right)\sigma + \frac{\bar{b}}{d}h\right). \quad (2.54)$$

Inserting this into the gauge-fixing term in (2.52), integrating by parts, rewriting in terms of Lichnerowicz Laplacians and using the York decomposition, we obtain the gauge-fixing term

$$S_{GF} = -\frac{Z_{GF}}{2a} \int d^d x \sqrt{g} \left[\hat{\xi}_\mu \left(\Delta_{L1} - \frac{2\bar{R}}{d} \right) \hat{\xi}^\mu + \frac{(d-1)^2}{d^2} \hat{\sigma} \left(\Delta_{L0} - \frac{\bar{R}}{d-1} \right) \hat{\sigma} + \frac{(d-1)\bar{b}}{d^2} \hat{\sigma} \sqrt{\Delta_{L0}} \sqrt{\Delta_{L0} - \frac{\bar{R}}{d-1}} h + \frac{\bar{b}^2}{d^2} h \Delta_{L0} h \right]. \quad (2.55)$$

We see that a specific combination of scalar degrees of freedom appears in the gauge-fixing term. It is sometimes convenient to write the scalar sector in terms of the gauge-invariant variable s defined in eq. (2.40) and this new degree of freedom which, in terms of the original fields, is given by

$$\chi = \sigma + \frac{b}{(d-1-b)\Delta_{L0} - \bar{R}} s = \frac{(d-1)\Delta_{L0} - \bar{R}}{(d-1-b)\Delta_{L0} - \bar{R}} \sigma + \frac{b(1+dm)}{(d-1-b)\Delta_{L0} - \bar{R}} h. \quad (2.56)$$

A short calculation shows that the Jacobian of the transformation $(\sigma, h) \rightarrow (s, \chi)$ is one.

In terms of the new variable, the gauge-fixing action is

$$S_{GF} = -\frac{Z_{GF}}{2a} \int d^d x \sqrt{g} \left[\xi_\mu \left(\Delta_{L1} - \frac{2\bar{R}}{d} \right)^2 \xi^\mu + \frac{(d-1-b)^2}{d^2} \chi \Delta_{L0} \left(\Delta_{L0} - \frac{\bar{R}}{d-1-b} \right)^2 \chi \right]. \quad (2.57)$$

From (2.56) we see that χ transforms in the same way as σ . Thus ξ and χ can be viewed as the gauge degrees of freedom and h^{TT} and s as the physical degrees of freedom. We will use this parametrization later.

The ghost action (2.52) for this gauge-fixing contains a non-minimal operator [1]. We also decompose the ghost into transverse and longitudinal parts

$$C_\nu = C_\nu^T + \bar{\nabla}_\nu C^L = C_\nu^T + \bar{\nabla}_\nu \frac{1}{\sqrt{-\nabla^2}} C'^L, \quad (2.58)$$

and the same for $\bar{C}_\mu$. (This change of variables has unit Jacobian). The ghost action then splits in two terms

$$S_{gh} = i \int d^d x \sqrt{g} \left[\bar{C}^{T\mu} \left(\Delta_{L1} - \frac{2\bar{R}}{d} \right) C_\mu^T + 2 \frac{d-1-b}{d} \bar{C}'^L \left(\Delta_{L0} - \frac{\bar{R}}{d-1-b} \right) C'^L \right]. \quad (2.59)$$

Note for future reference that the gauge-fixing condition and the ghost action are pathological for $b = d - 1$.

Let us observe that Z_{GF} appears in the combination Z_{GF}/a , where a is a dimensionless gauge parameter, while Z_{GF} is a constant of dimension $d-2$. There are then two natural options regarding the constant Z_{GF} . The first choice is to treat it as a fixed parameter (later we shall identify it with a power of the cutoff). This leads to simpler formulas for many expressions, and we shall use it extensively later. It is however not appropriate for the discussion of perturbative Einstein gravity. The reason is that in the limit $G \rightarrow 0$ the coefficient Z_N of the Hessian diverges. If we keep Z_{GF} and a constant in the limit, then the gauge-fixing term becomes negligible relative to the rest of the quadratic action. The gauge fluctuations remain unsuppressed and one can anticipate divergences. This is exactly what happens, as we shall mention later on. One can compensate the behavior of Z_N by keeping Z_{GF} fixed and letting simultaneously $a \rightarrow 0$.

Alternatively, one can set $Z_{GF} = Z_N$. In this case, in the Gaussian limit $G \rightarrow 0$, the kinetic terms of the gauge-invariant and gauge degrees of freedom scale in the same way and one obtains sensible results for all values of a and b .

In the following we shall discuss also different choices for the gauge fixing. One is the so-called “unimodular physical gauge”, where one sets $\hat{\xi}_\mu = 0$ and $h = 0$. As shown in [21], this is equivalent to the above standard gauge in the limit $b \rightarrow \pm\infty$ and $a \rightarrow 0$.

2.7 Four-derivative gauge-fixing terms

In HDG it is customary to use gauge-fixing terms that contain four derivatives. In order to further check gauge-independence of the results, in section 3.2 we consider the gauge fixing in our previous paper [19] but now with arbitrary gauge parameters.

We can simply repeat the same procedure to derive the gauge-fixing and FP terms but with additional higher derivative operator

$$Y_{\mu\nu} \equiv \bar{g}_{\mu\nu} \square + c \bar{\nabla}_\mu \bar{\nabla}_\nu - f \bar{\nabla}_\nu \bar{\nabla}_\mu, \quad (2.60)$$

where c and f are additional gauge parameters:

$$\begin{aligned} \mathcal{L}_{GF+FP}/\sqrt{g} &= i\delta_B \left[\bar{C}_\mu Y^{\mu\nu} \left(F_\nu - \frac{a}{2Z_{GF}} B_\nu \right) \right] / \delta\lambda \\ &= -\frac{a}{2Z_{GF}} \tilde{B}_\mu Y^{\mu\nu} \tilde{B}_\nu + \frac{Z_{GF}}{2a} F_\mu Y^{\mu\nu} F_\nu + i\bar{C}_\mu Y^{\mu\nu} \Delta_{\mu\nu}^{(gh)} C^\nu, \end{aligned} \quad (2.61)$$

where the gauge-fixing function F_μ is defined in (2.47), and $\tilde{B}_\mu$ and $\Delta_{\mu\nu}^{(gh)}$ in (2.53). Here we should notice that the dimension of the constant Z_{GF} is changed to $d-4$.

Note also that the field B_μ was an auxiliary field in the two-derivative gauge-fixing case (2.52), but here it is dynamical. With higher derivative gauge fixing, quite often only the contribution $\Delta_{\mu\nu}^{(gh)}$ is incorporated but that from $Y^{\mu\nu}$ in the FP ghost kinetic term is ignored, and then it is claimed that somehow the contribution from the “third ghost” $-\frac{1}{2} \log(\det(Y_{\mu\nu}))$ must be added. We see here that this is automatic in the BRST invariant formulation, because we have contributions $-\log(\det(Y^{\mu\nu}))$ from the FP ghost kinetic term and $\frac{1}{2} \log(\det(Y^{\mu\nu}))$ from the field B_μ , giving the same result.

Using the York decomposition (2.24), we can calculate the contributions without fixing gauge parameters and so can check the gauge dependence directly here. We find that the terms (2.61) are cast into

$$\begin{aligned} & -\frac{a}{2Z_{GF}} \left[B_\mu^T \left(\Delta_{L1} - \frac{1-f}{d} \bar{R} \right) B^{\mu T} + B^L \Delta_{L0} \left\{ (1+c-f) \Delta_{L0} - \frac{1-f}{d} \bar{R} \right\} B^L \right] \\ & + \frac{Z_{GF}}{2a} \left[\xi_\mu \left(\Delta_{L1} - \frac{2}{d} \bar{R} \right)^2 \left(\Delta_{L1} - \frac{1-f}{d} \bar{R} \right) \xi^\mu \right. \\ & \quad \left. + \left(\frac{d-1-b}{d} \right)^2 \chi \Delta_{L0} \left(\Delta_{L0} - \frac{\bar{R}}{d-1-b} \right)^2 \left\{ (1+c-f) \Delta_{L0} - \frac{1-f}{d} \bar{R} \right\} \chi \right] \\ & + i\bar{C}_\mu^T \left(\Delta_{L1} - \frac{1-f}{d} \bar{R} \right) \left(\Delta_{L1} - \frac{2}{d} \bar{R} \right) C^{\mu T} \\ & + 2i \frac{d-1-b}{d} \bar{C}^L \Delta_{L0} \left(\Delta_{L0} - \frac{\bar{R}}{d-1-b} \right) \left\{ (1+c-f) \Delta_{L0} - \frac{1-f}{d} \bar{R} \right\} C^L, \end{aligned} \quad (2.62)$$

where we have defined transverse and longitudinal parts of the $\tilde{B}_\mu$ and FP ghosts $\bar{C}_\mu$ and C_μ , as usual.

In section 3.1 we will discuss the gauge-independence of the theory in the general quadratic gauge. In section 4.5 we will restrict ourselves to the class of gauges where $c = f = 1$ but with generic a, b . This is equivalent to inserting a Lichnerowicz Laplacian Δ_{L1} in the quadratic gauge fixing term (2.52), or in other words to set $Y^{\mu\nu} = \bar{g}^{\mu\nu} \Delta_{L1}$. After performing the York decomposition, this yields additional factors Δ_{L1} and Δ_{L0} in the quadratic actions of ξ_μ and χ , equation (2.57). The resulting additional determinants are offset by the determinant of the operator coming from the B_μ sector, as will become clear in the following.

3 Universality in four-derivative gravity in $d = 4$

In this section we consider the theory with $Z_N = 0$ in $d = 4$ on an Einstein background. Since the equation of motion is automatically satisfied, one would expect the effective action to be gauge- and parametrization-independent. We will check that this is indeed the case at the formal level of determinants. In the following section we will have a more explicit check of this property in the expressions for the divergences.

3.1 General two-derivative gauge fixing

The Hessian of pure higher-derivative gravity in $d = 4$ has the simple form (2.45), independent of the choice of parametrization. It is expressed entirely in terms of the gauge-invariant variables h^{TT} and s . These fields do not appear in the gauge-fixing action, so their contribution to the one-loop effective action is given by:

$$\text{Det}\left(\Delta_{L2} - \frac{\bar{R}}{2}\right)^{-1/2} \text{Det}\left(\Delta_{L2} - \left(1 + \frac{2\alpha}{\beta}\right)\bar{R}\right)^{-1/2} \text{Det}\Delta_{L0}^{-1/2} \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3}\right)^{-1/2}. \quad (3.1)$$

We choose the gauge as in section 2.6. It is convenient to express the gauge fixing term as in (2.57). The fields ξ and χ contribute to the one-loop action the terms (as ordered):

$$\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right)^{-1} \text{Det}\Delta_{L0}^{-1/2} \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3-b}\right)^{-1}. \quad (3.2)$$

The ghost action (2.59) gives the determinants

$$\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right) \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3-b}\right), \quad (3.3)$$

and finally the Jacobians of the change of variables from $h_{\mu\nu}$ to $h_{\mu\nu}^{TT}, \xi^\mu, \sigma$ and h is:

$$\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right)^{1/2} \text{Det}\Delta_{L0}^{1/2} \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3}\right)^{1/2}. \quad (3.4)$$

Note that the gauge parameter a only appears in the Hessian in the prefactors and not in the operators. The only dependence on the gauge parameter b is in two determinants that cancel. This proves the gauge independence of the one-loop action. The final result is

$$\frac{\sqrt{\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right)}}{\sqrt{\text{Det}\left(\Delta_{L2} - \frac{\bar{R}}{2}\right)} \sqrt{\text{Det}\left(\Delta_{L2} - \left(1 + \frac{2\alpha}{\beta}\right)\bar{R}\right)} \sqrt{\text{Det}\Delta_{L0}}} . \quad (3.5)$$

3.2 General four-derivative gauge fixing

If we use the general four-derivative gauge fixing of section 2.7, we have again the determinants (3.1) coming from the physical degrees of freedom and (3.4) coming from the Jacobians. The determinants coming from the fields ξ_μ and χ are

$$\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right)^{-1} \text{Det}\left(\Delta_{L1} - \frac{1-f}{4}\bar{R}\right)^{-1/2} \quad (3.6)$$

$$\times \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3-b}\right)^{-1} \text{Det}\left(\Delta_{L0} - \frac{1-f}{4(1+c-f)}\bar{R}\right)^{-1/2}, \quad (3.7)$$

the ghosts give

$$\begin{aligned} & \text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right) \text{Det}\left(\Delta_{L1} - \frac{1-f}{4}\bar{R}\right) \\ & \times \text{Det}\Delta_{L0} \text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3-b}\right) \text{Det}\left(\Delta_{L0} - \frac{1-f}{4(1+c-f)}\bar{R}\right), \end{aligned} \quad (3.8)$$

and finally we have the contributions of the B_μ field

$$\text{Det}\left(\Delta_{L1} - \frac{1-f}{4}\bar{R}\right)^{-1/2} \text{Det}\Delta_{L0}^{-1/2} \text{Det}\left(\Delta_{L0} - \frac{1-f}{4(1+c-f)}\bar{R}\right)^{-1/2}. \quad (3.9)$$

Putting everything together, we see that all the terms depending on the gauge parameters cancel out and the remaining ones give again (3.5). Thus we have explicitly shown, at this formal level, that the results are completely gauge independent.

3.3 Physical gauge

In the calculation of sections 3.1, and even more in section 3.2, there is a large number of cancellations between various determinants. Consider instead the “physical” gauge $\hat{\xi}_\mu = 0$, $h = 0$, discussed in [20, 21]. It leaves only the fields h^{TT} and $\hat{\sigma}$, and no Jacobians. The Hessians of h^{TT} and $\hat{\sigma}$ are given by (2.35) and (2.37) respectively. There are a real scalar ghost and a real transverse vector ghost, with ghost operators Δ_{L0} and $\Delta_{L1} - R/2$. Putting together these terms, one immediately obtains the effective action (3.5). In fact this is the most direct way of getting it, because there is no cancellation of determinants between unphysical degrees of freedom and ghosts.

3.4 The conformal case

Now we consider the conformal case where $\beta/\alpha = -3$. The effective action in this case cannot be simply obtained as a particular case of eq.(3.5), because the action is invariant also under Weyl transformations and this requires a separate gauge fixing.

The Hessian is only nonzero in the spin-two sector and is given by (2.46). For the Weyl invariance we can gauge fix $h = 0$, without any ghost because h transforms under Weyl transformations by a shift. For diffeomorphisms we choose a standard gauge fixing of the form (2.55). Since $h = 0$, the value of b is immaterial. Equation (2.56) is replaced by $\chi = \sigma$, so the decomposition of the gauge-fixing and ghost actions, and the corresponding determinants, are the same as in section 2.6, with $b = 0$.

We use the fields $\hat{\xi}$ and $\hat{\sigma}$ (see (2.25)) in such a way that no Jacobian is needed. The one-loop effective action is

$$\frac{\sqrt{\text{Det}\left(\Delta_{L1} - \frac{\bar{R}}{2}\right)}\sqrt{\text{Det}\left(\Delta_{L0} - \frac{\bar{R}}{3}\right)}}{\sqrt{\text{Det}\left(\Delta_{L2} - \frac{\bar{R}}{2}\right)}\sqrt{\text{Det}\left(\Delta_{L2} - \frac{\bar{R}}{3}\right)}}. \quad (3.10)$$

The two terms in the denominator come from the TT part. The terms in the numerator come from the ghosts (power 1) and from the gauge fixing term (power $-1/2$). Note that there is no dependence on the gauge parameter a . It only appears in the prefactors of the quadratic action, which drop out.

Alternatively we can choose a different gauge. For Weyl transformations we still choose $h = 0$, which leaves no ghost term. For diffeomorphisms we choose the second type of physical gauge explained in the end of section III.B of [21], namely $\hat{\sigma} = 0$ and $\hat{\xi} = 0$. This is equivalent to taking the Landau gauge limit $a \rightarrow 0$. The ghosts are a real scalar and a real transverse vector and the ghost operators are $\Delta_{L0} - \bar{R}/3$ and $\Delta_{L1} - \bar{R}/2$. The effective action is given again by (3.10).

4 The divergences

4.1 Derivation

The one-loop effective action contains a divergent part

$$\begin{aligned} \Gamma_k = & \int d^d x \sqrt{g} \left[\frac{A_1}{16\pi d} k^d + \frac{B_1}{16\pi(d-2)} k^{d-2} \bar{R} \right. \\ & \left. + \beta_\alpha \frac{k^{d-4}}{d-4} \bar{R}^2 + \beta_\beta \frac{k^{d-4}}{d-4} \bar{R}_{\mu\nu} \bar{R}^{\mu\nu} + \beta_\gamma \frac{k^{d-4}}{d-4} \bar{R}_{\mu\nu\rho\sigma} \bar{R}^{\mu\nu\rho\sigma} \right], \end{aligned} \quad (4.1)$$

where k stands for a cutoff. In $d = 4$, the power divergences $k^{d-4}/(d-4)$ of the last three terms are replaced by terms $\log(k/\mu)$, where we introduced a reference mass scale μ . In general one would have separate Riemann squared, Ricci squared and R^2 terms, but on an Einstein space (2.8) the latter two merge into a single term proportional to $\bar{R}^2$. Then, we will denote

$$C_1 = \beta_\alpha + \frac{1}{d} \beta_\beta, \quad (4.2)$$

the coefficient of the $\bar{R}^2$ divergence and $D_1 = \beta_\gamma$ the coefficient of the $\bar{R}_{\mu\nu\rho\sigma} \bar{R}^{\mu\nu\rho\sigma}$ divergence. We note that in I we used a maximally symmetric background, where $\bar{R}_{\mu\nu\rho\sigma} \bar{R}^{\mu\nu\rho\sigma} = \frac{2}{d(d-1)} \bar{R}^2$. Thus, the coefficient C_1 of I corresponds to the combination $C_1 + \frac{2}{d(d-1)} D_1$.

We describe here the algorithm that we use to derive the coefficients. Instead of Γ_k we shall evaluate the derivative [23, 24]:

$$\dot{\Gamma}_k = \int d^d x \sqrt{g} \left[\frac{A_1}{16\pi} k^d + \frac{B_1}{16\pi} k^{d-2} \bar{R} + (\beta_\alpha \bar{R}^2 + \beta_\beta \bar{R}_{\mu\nu} \bar{R}^{\mu\nu} + \beta_\gamma \bar{R}_{\mu\nu\rho\sigma} \bar{R}^{\mu\nu\rho\sigma}) k^{d-4} + \dots \right], \quad (4.3)$$

where the overdot stands for $k \frac{d}{dk}$. The advantage of this procedure is that this quantity is convergent. Independently of the renormalizability properties of the theory, it defines a flow on the space of all couplings that are permitted by the symmetries of the system. The divergences of (4.1) in the limit $k \rightarrow \infty$ are then found by integrating the differential equation (4.3) from

some initial scale k_0 up to k . The coefficients A_1 , B_1 , β_α , β_β and β_γ enter in the beta functions of the couplings Λ , G , α , β , γ , but we postpone a discussion of this point to section 7.

The technique employed to evaluate the right hand side of (4.3) is similar to the one described in I, but this time the cutoff is taken to be a function of the Lichnerowicz Laplacians instead of the Bochner Laplacian $-\bar{\nabla}^2$, as in I. In each spin sector, the Lichnerowicz operator Δ_L is replaced by the regularized one $P_k(\Delta_L) = \Delta_L + R_k(\Delta_L)$, where we use the optimized cutoff $R_k(\Delta_L) = (k^2 - \Delta_L)\theta(k^2 - \Delta_L)$. Then $\dot{\Gamma}_k$ is given by

$$\dot{\Gamma}_k = \frac{1}{2}\text{Tr} \left(\frac{\dot{\Delta}_k^{(2)}}{\Delta_k^{(2)}} \right) + \frac{1}{2}\text{Tr} \left(\frac{\dot{\Delta}_k^{(1)}}{\Delta_k^{(1)}} \right) + \frac{1}{2}\text{Tr} \left(\frac{\dot{\Delta}_k^{(0)}}{\Delta_k^{(0)}} \right) - \text{Tr} \left(\frac{\dot{\Delta}_{gh,k}^{(1)}}{\Delta_{gh,k}^{(1)}} \right) - \text{Tr} \left(\frac{\dot{\Delta}_{gh,k}^{(0)}}{\Delta_{gh,k}^{(0)}} \right), \quad (4.4)$$

where Δ_k are the kinetic operators that appear in the each spin sector and the numerator is $\dot{\Delta}_k = \dot{R}_k(\Delta_L) = 2k^2\theta(k^2 - \Delta_L)$. The step function cuts off the trace to eigenvalues of Δ_L that are smaller than k^2 . For example in the spin-two sector

$$\begin{aligned} \frac{1}{2}\text{Tr} \left(\frac{\dot{\Delta}_k^{(2)}}{\Delta_k^{(2)}} \right) &= \frac{1}{2} \frac{1}{(4\pi)^{d/2}} \left[W(\Delta_L^{(2)}, 0) \left(Q_{d/2} b_0(\Delta_L^{(2)}) + Q_{d/2-1} b_2(\Delta_L^{(2)}) + Q_{d/2-2} b_4(\Delta_L^{(2)}) \right) \right. \\ &\quad + W'(\Delta_L^{(2)}, 0) \bar{R} \left(Q_{d/2} b_0(\Delta_L^{(2)}) + Q_{d/2-1} b_2(\Delta_L^{(2)}) \right) \\ &\quad \left. + \frac{1}{2} W''(\Delta_L^{(2)}, 0) \bar{R}^2 \left(Q_{d/2} b_0(\Delta_L^{(2)}) \right) + \dots \right], \end{aligned} \quad (4.5)$$

where $W(\Delta_L^{(2)}, \bar{R}) = \frac{\dot{\Delta}_{Lk}^{(2)}}{\Delta_{Lk}^{(2)}}$ and primes denote derivatives with respect to $\bar{R}$. The coefficients Q_n and the heat kernel coefficients b_n are listed in Appendix A.

Similar formulas hold for the spin one and spin zero sectors and for the ghosts. In the scalar term, $\Delta_k^{(0)}$ is a two-by-two matrix, and the fraction has to be understood as the product of $\dot{\Delta}_k^{(0)}$ with the inverse of $\Delta_k^{(0)}$. The functional trace thus involves also a trace over this two-by-two matrix. With these data one can write the expansion of (4.4) in powers of $\bar{R}$, and comparing with (4.3) one can read off the coefficients A_1 , B_1 and C_1 .

4.2 Results

From now on we will deal with dimensionless variables

$$\tilde{\alpha} = k^{4-d}\alpha, \quad \tilde{\beta} = k^{4-d}\beta, \quad \tilde{\Lambda} = k^{-2}\Lambda, \quad \tilde{G} = k^{d-2}G, \quad (\tilde{Z}_N = k^{2-d}Z_N), \quad (4.6)$$

and

$$\tilde{Z}_{GF} = \begin{cases} k^{2-d}Z_{GF} & \text{for two-derivative gauge fixing (2.52)} \\ k^{4-d}Z_{GF} & \text{for four-derivative gauge fixing (2.61)} \end{cases}. \quad (4.7)$$

In I, the couplings $\tilde{\alpha}$ and $\tilde{\beta}$ were absent, and $\tilde{G}$ only appeared in the Hessian through the overall prefactor Z_N . Since this prefactor cancelled between numerator and denominator in (4.4), the divergences were independent of $\tilde{G}$. This does no longer happen in the present theory, so the coefficients A_1 , B_1 and C_1 depend in general on the dimension d , on the measure parameters m and ω , on the gauge parameters a and b , and on the couplings $\tilde{\Lambda}$, $\tilde{G}$, $\tilde{\alpha}$ and $\tilde{\beta}$. The general expressions for the coefficients of the divergences are extremely complicated and not instructive. We will therefore only write them in certain limits where they simplify enough.

Compared to I, we have to take into account the additional dependence on the couplings $\tilde{G}$, $\tilde{\alpha}$ and $\tilde{\beta}$. We will consider two limiting situations. One is the limit $\tilde{\alpha} \rightarrow 0$ and $\tilde{\beta} \rightarrow 0$. In this case the beta functions should reduce to those of Einstein-Hilbert gravity and match with those of I (modulo scheme dependences, due to the different form of the cutoff in this paper). Still, the present results are more general because we consider a generic Einstein background, which allows us to distinguish at least two of the higher derivative couplings, whereas in I a maximally symmetric background was used, allowing us to calculate the coefficient of a single combination of the higher derivative couplings. We will refer to this as the “Einstein-Hilbert limit” (EH limit).²

The other limit consists in taking $\tilde{Z}_N \rightarrow 0$, which is equivalent to $\tilde{G} \rightarrow \infty$. In this case one is left with functions of the higher derivative couplings only. We will refer to this as the “four-derivative gravity limit” (4DG-limit).

There is still the freedom of choosing between the two-derivative and four-derivative gauges. In sections 4.3 and 4.4 we will consider the 4DG- and the EH-limits of the theory, using the two-parameter family of two-derivative gauges introduced in section 2.6. In section 4.5 we will discuss the changes that occur when using the two-parameter family of four-derivative gauges introduced in section 2.7. It will turn out that in order to take the EH- and 4DG-limits, different choices have to be made regarding the overall gauge-fixing coefficient Z_{GF} . These are spelled out in detail in the following sections.

4.3 The 4DG-limit ($\tilde{Z}_N \rightarrow 0$)

For this calculation we set $\tilde{Z}_{GF} = 1$. The coefficients A_1 , B_1 and D_1 turn out to be universal (i.e. independent of the gauge and parametrization) in any dimension:

$$\begin{aligned} A_1 &= \frac{8(d-2)}{(4\pi)^{d/2-1}\Gamma(d/2)}, \\ B_1 &= \frac{2}{(4\pi)^{d/2-1}3d^2\tilde{\beta}(4(d-1)\tilde{\alpha}+d\tilde{\beta})\Gamma(d/2)}[48d(d^3-2d^2-d+2)\tilde{\alpha}^2 \\ &\quad +4(d^5-12d^4+41d^3-102d^2+36d+48)\tilde{\alpha}\tilde{\beta}+(d^5-14d^4+30d^3-60d^2-72d+96)\tilde{\beta}^2], \\ D_1 &= \frac{-1050+589d-34d^2+d^3}{(4\pi)^{d/2}360\Gamma(d/2)}. \end{aligned} \tag{4.8}$$

The coefficient C_1 has a complicated dependence on m , ω , a , b , $\tilde{\alpha}$ and $\tilde{\beta}$, that we do not report here. A special case will be given below in (4.21). However, in $d = 4$ all four coefficients, including C_1 , are universal:

$$\begin{aligned} A_1 &= \frac{4}{\pi}, \\ B_1 &= \frac{15\tilde{\alpha}-14\tilde{\beta}}{3\pi\tilde{\beta}}, \\ C_1 &= \frac{1200\tilde{\alpha}^2+200\tilde{\alpha}\tilde{\beta}-183\tilde{\beta}^2}{1920\pi^2\tilde{\beta}^2}, \\ D_1 &= \frac{413}{2880\pi^2}. \end{aligned} \tag{4.9}$$

²The same results can be obtained, at least in some cases, by taking the limit $\tilde{Z}_N \rightarrow \infty$, which is the same as considering the perturbative regime $\tilde{G} \rightarrow 0$.

This is an explicit consequence of the statement made in section 3.1, that the one-loop effective action of higher derivative gravity is independent of the parametrization and gauge choice in $d = 4$.

The expressions for C_1 and D_1 given above are in agreement with the standard results for the beta functions in HDG [8, 26] (also derived by means of (4.4) in [27, 28, 19]):

$$\begin{aligned}\beta_{\tilde{\alpha}} &= \frac{1}{(4\pi)^2} \frac{90\tilde{\alpha}^2 - 23\tilde{\beta}^2 - 338\tilde{\gamma}^2 + 15\tilde{\alpha}\tilde{\beta} - 120\tilde{\alpha}\tilde{\gamma} - 199\tilde{\beta}\tilde{\gamma}}{9(\tilde{\beta} + 4\tilde{\gamma})^2}, \\ \beta_{\tilde{\beta}} &= \frac{1}{(4\pi)^2} \frac{371}{90}, \\ \beta_{\tilde{\gamma}} &= \frac{1}{(4\pi)^2} \frac{413}{180}.\end{aligned}\tag{4.10}$$

Let us consider what happens if we choose $\tilde{Z}_{GF} = \tilde{Z}_N$, as is usually done in Einstein-Hilbert theory. In the 4DG-limit, the coefficients A_1 and D_1 are still given by the formulas given above, but B_1 and C_1 are different, and gauge-dependent. However, for the exponential parametrization ($\omega = \frac{1}{2}$) in $d = 4$, one gets m -independent result

$$\begin{aligned}B_1 &= \frac{15\tilde{\alpha}(b-3)^2 + \tilde{\beta}(3a - 14(b-3)^2)}{3\pi(b-3)^2\tilde{\beta}}, \\ C_1 &= \frac{\tilde{\beta}^2(240a^2 + 40a(b^2 - 18b + 45) - 183(b-3)^4) + 1200\tilde{\alpha}^2(b-3)^4 + 200\tilde{\alpha}\tilde{\beta}(b-3)^4}{1920\pi^2(b-3)^4\tilde{\beta}^2}.\end{aligned}\tag{4.11}$$

We note that in the limit $a \rightarrow 0$ these reproduce the universal formulas given above. This observation agrees with the discussion in the end of section 2.6. In the 4DG-limit, at fixed k , one sets $\tilde{Z}_N = 0$. For fixed a and k , and with the choice $\tilde{Z}_{GF} = \tilde{Z}_N$, this implies that the gauge fixing term vanishes too. The situation can be fixed by letting $a \rightarrow 0$, with the ratio $\tilde{Z}_{GF}/a$ fixed and finite. In conclusion, the gauge-dependence that occurs in (4.11) for $a \neq 0$ is the artifact of a bad gauge-fixing procedure. The correct result is given by (4.9).

This problem does not arise in the standard calculation of the beta functions of HDG in $d = 4$, because there the gauge fixing is of the type considered in section 3.2. When the gauge-fixing term contains four derivatives, its overall coefficient is dimensionless in $d = 4$ and there is never the temptation to introduce a factor $\tilde{Z}_N$.

4.4 The EH limit

Let us now consider the EH limit $\tilde{\alpha}, \tilde{\beta} \rightarrow 0$. In this case it is not appropriate to set $\tilde{Z}_{GF} = 1$, as we did in the preceding subsection, because of the issue in the gauge fixing discussed in the end of section 2.6. Indeed if we insisted with this choice we would find that for $\tilde{\Lambda} = 0$ only A_1 and D_1 are universal:

$$\begin{aligned}A_1 &= \frac{4(d-3)}{(4\pi)^{d/2-1}\Gamma(d/2)}, \\ D_1 &= \frac{d^3 - 35d^2 + 606d - 1080}{(4\pi)^{d/2}720\Gamma(d/2)}.\end{aligned}\tag{4.12}$$

whereas B_1 and C_1 would contain $\tilde{G}$ and diverge for $\tilde{G} \rightarrow 0$. For this reason we switch now to the choice $\tilde{Z}_{GF} = \tilde{Z}_N$, which is more appropriate to discuss this limit.

We find that A_1 and D_1 are still given by (4.12). In $d = 4$ we have:

$$\begin{aligned}
A_1 &= \frac{1}{\pi}, \\
B_1 &= \frac{1}{12\pi(b-3)^2(4m+1)} \left[3a(-18b(4m+1)^2(2\omega-1) + 3(2\omega-1)(4bm+b)^2 \right. \\
&\quad + 8m(66(2m+1)\omega - 66m - 31) + 70\omega - 31 + b^2(4m+1)(72m\omega - 36m + 18\omega - 47) \\
&\quad \left. - 6b(4m+1)(30(4m+1)\omega - 60m - 53) + 9(48m(10m\omega - 5m + 5\omega - 6) + 26\omega - 55) \right], \\
D_1 &= \frac{53}{720\pi^2}.
\end{aligned} \tag{4.13}$$

We note that $\tilde{G}$ does not appear in these expressions, so that they have a smooth limit $\tilde{G} \rightarrow 0$. The expression for C_1 is still too long to be written but is likewise independent of $\tilde{G}$. We give it only for two cases: linear parametrization $m = \omega = 0$

$$\begin{aligned}
C_1 &= \frac{1}{1920\pi^2(b-3)^4} [15a^2(3b^4 - 36b^3 + 162b^2 - 324b + 259) \\
&\quad - 20a(3b^4 - 36b^3 + 176b^2 - 360b + 297) + 24(7b^4 - 59b^3 + 223b^2 - 381b + 252)].
\end{aligned} \tag{4.14}$$

and exponential parametrization $\omega = 1/2$, in which case the result is automatically independent of m :

$$C_1 = \frac{240a^2 + 40a(b^2 - 18b + 9) - 3(29b^4 - 348b^3 + 1526b^2 - 3372b + 2709)}{1920\pi^2(b-3)^4}. \tag{4.15}$$

One can plot these functions for fixed parametrization or for fixed gauge. One obtains plots that are very similar to those shown in I. They will not be repeated here.

We report for completeness the expressions B_1 and C_1 for $d = 4$ in the “unimodular physical” gauge $b \rightarrow \pm\infty$, $a \rightarrow 0$, for $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{G}$ all tending to zero:

$$\begin{aligned}
B_1 &= \frac{36m(2\omega-1) + 18\omega - 47}{12\pi}, \\
C_1 &= \frac{180m^2(1-2\omega)^2 + 10m(36\omega^2 - 44\omega + 13) + 45\omega^2 - 65\omega + 14}{160\pi^2}.
\end{aligned} \tag{4.16}$$

4.5 Results with the four-derivative gauge fixing

In the preceding sections we used the two-derivative gauge-fixing terms introduced in section 2.6. We now discuss briefly the results when the four-derivative gauge-fixing terms of section 2.7 are used instead.

The formal discussion of section 3 indicates that in the 4DG-limit in four dimensions, the divergences should be universal. Explicit calculation confirms that in this case the coefficients A_1 , B_1 , C_1 , D_1 are indeed given again by (4.9). Furthermore, the coefficients A_1 , B_1 and D_1 are universal in any dimension and agree with those given in (4.8).

In the EH limit the coefficients A_1 , and D_1 in $d = 4$ are again as in (4.13), but the others are slightly different. For B_1 one has in general

$$\begin{aligned}
B_1 &= \frac{1}{12\pi(b-3)^2(4m+1)} \left[6a(-18b(4m+1)^2(2\omega-1) + 3(2\omega-1)(4bm+b)^2 \right. \\
&\quad + 8m(66(2m+1)\omega - 66m - 31) + 70\omega - 31 + b^2(4m+1)(72m\omega - 36m + 18\omega - 47) \\
&\quad \left. - 6b(4m+1)(30(4m+1)\omega - 60m - 53) + 9(48m(10m\omega - 5m + 5\omega - 6) + 26\omega - 55) \right].
\end{aligned} \tag{4.17}$$

Similarly the coefficient C_1 in the linear parametrization $m = \omega = 0$ is

$$C_1 = \frac{1}{1920\pi^2(b-3)^4} [30a^2(3b^4 - 36b^3 + 162b^2 - 324b + 259) - 5a(15b^4 - 180b^3 + 898b^2 - 1860b + 1575) + 24(7b^4 - 59b^3 + 223b^2 - 381b + 252)] . \quad (4.18)$$

and in the exponential parametrization $\omega = 1/2$:

$$C_1 = \frac{480a^2 + 80a(b^2 - 15b + 9) - 3(29b^4 - 348b^3 + 1526b^2 - 3372b + 2709)}{1920\pi^2(b-3)^4} . \quad (4.19)$$

The result in the unimodular physical gauge $b \rightarrow \pm\infty$, $a \rightarrow 0$ agrees with (4.16).

4.6 Exponential parametrization and unimodular gauge

It was found in I that choosing the exponential parametrization $\omega = 1/2$ and the unimodular gauge $b \rightarrow \pm\infty$, all dependence on the other two parameters m and a , as well as the dependence on the cosmological constant $\tilde{\Lambda}$, drop out.

This result holds true also in the present context, in any dimension and independently of whether the gauge fixing contains two or four derivatives and independently of the treatment of the constant $\tilde{Z}_{GF}$. The resulting coefficients are still too cumbersome to write, so we give them again only in two limits.

In the 4DG-limit the coefficients A_1 , B_1 and D_1 are given by (4.8) and

$$C_1 = \frac{1}{720(4\pi)^{d/2}d^3\tilde{\beta}^2(4(d-1)\tilde{\alpha} + d\tilde{\beta})^2\Gamma(d/2)} \left[46080\tilde{\alpha}^4(d-1)^2d^2(d^2-d-2) + 1920\tilde{\alpha}^3\tilde{\beta}d(d^6-3d^5+85d^4-285d^3+106d^2+288d-192) + 16\tilde{\alpha}^2\tilde{\beta}^2(5d^8-92d^7+917d^6-1886d^5+7964d^4-49268d^3+42360d^2+25920d-23040) + 8\tilde{\alpha}\tilde{\beta}^3d(5d^7-132d^6+1235d^5-4716d^4+10988d^3-28800d^2-14880d+46080) + \tilde{\beta}^4(5d^8-142d^7+1288d^6-4628d^5+7560d^4-19200d^3-17280d^2-46080d+92160) \right] . \quad (4.20)$$

In the EH limit we have

$$\begin{aligned} A_1 &= \frac{4(d-3)}{(4\pi)^{d/2-1}\Gamma(d/2)}, \\ B_1 &= \frac{(d^3-15d^2+24d-72)}{3d(4\pi)^{d/2-1}\Gamma(d/2)}, \\ C_1 &= \frac{5d^5-147d^4+1240d^3-3612d^2+2880d-17280}{1440d^2(4\pi)^{d/2}\Gamma(d/2)} \\ D_1 &= \frac{d^3-35d^2+606d-1080}{720(4\pi)^{d/2}\Gamma(d/2)} . \end{aligned} \quad (4.21)$$

We note that the formulas for B_1 and $C_1 + \frac{2}{d(d-1)}D_1$ do not agree with equation (5.7) in I. This is due to the different choice of cutoff (a function of the Bochner Laplacian in I and of the Lichnerowicz Laplacians here). Specializing to $d = 4$, the formula for $C_1 + \frac{2}{d(d-1)}D_1$ agrees with (5.6) in I but B_1 does not. This is a signal of the lack of universality of this coefficient.

5 Conformal gravity in $d = 4$

Independently of the choice of gauge and parametrization, the effective action for conformal gravity is given formally by (3.10). This leads to the following flow for the effective action

$$\dot{\Gamma}_k = \frac{1}{2} \text{Tr} \left(\frac{\dot{\Delta}_k^{(2)}}{\Delta_k^{(2)}} \right) - \frac{1}{2} \text{Tr} \left(\frac{\dot{\Delta}_{gh,k}^{(1)}}{\Delta_{gh,k}^{(1)}} \right) - \frac{1}{2} \text{Tr} \left(\frac{\dot{\Delta}_{gh,k}^{(0)}}{\Delta_{gh,k}^{(0)}} \right), \quad (5.1)$$

where

$$\begin{aligned} \Delta_k^{(2)} &= \left(P_k(\Delta_{L2}) - \frac{1}{2} \bar{R} \right) \left(P_k(\Delta_{L2}) - \frac{1}{3} \bar{R} \right), \\ \Delta_k^{(1)} &= P_k(\Delta_{L1}) - \frac{1}{2} \bar{R}, \\ \Delta_k^{(0)} &= P_k(\Delta_{L0}) - \frac{1}{3} \bar{R}. \end{aligned} \quad (5.2)$$

Proceeding as before we find

$$A_1 = \frac{3}{(4\pi)^2}, \quad (5.3)$$

$$B_1 = \frac{41}{6(4\pi)^2}, \quad (5.4)$$

$$C_1 = -\frac{199}{180(4\pi)^2}, \quad (5.5)$$

$$D_1 = \frac{137}{60(4\pi)^2}. \quad (5.6)$$

We observe that the logarithmic divergences are given by

$$\frac{1}{(4\pi)^2} \left[b_4 \left(\Delta_{L2} - \frac{1}{2} \bar{R} \right) + b_4 \left(\Delta_{L2} - \frac{1}{3} \bar{R} \right) - b_4 \left(\Delta_{L1} - \frac{1}{2} \bar{R} \right) - b_4 \left(\Delta_{L2} - \frac{1}{2} \bar{R} \right) \right]. \quad (5.7)$$

The coefficients C_1 and D_1 agree with previous calculations reported in [29] and [30].

6 Duality

One of the main results of I was the existence of a discrete idempotent transformation leaving the functional measure, and the divergences, invariant. That invariance extends also to HDG, at least on an Einstein background. In any dimension and in any gauge we find that the divergences A_1, B_1, C_1 and D_1 have the duality symmetry:

$$\begin{aligned} A_1(\omega, m) &= A_1 \left(1 - \omega, -m - \frac{2}{d} \right), \\ B_1(\omega, m) &= B_1 \left(1 - \omega, -m - \frac{2}{d} \right), \\ C_1(\omega, m) &= B_1 \left(1 - \omega, -m - \frac{2}{d} \right), \\ D_1(\omega, m) &= C_1 \left(1 - \omega, -m - \frac{2}{d} \right). \end{aligned} \quad (6.1)$$

This duality is only manifest if the coefficient b enters in the gauge fixing (2.47) through the combination $\bar{b} = b(1 + dm)$. With other definitions of this coefficient a form of duality will still be present but it will have a much more complicated form.

The calculations reported here indicate that this duality is not limited to Einstein gravity. It will be interesting to understand more generally under what circumstances this property holds.

7 Concluding remarks

In this paper we have extended the analysis of I [1] from Einstein-Hilbert gravity to higher-derivative gravity containing the squares of the Ricci scalar and Ricci tensor. In four-dimensions the analysis is essentially complete, because the remaining independent invariant is a total derivative. In higher dimensions this is not so. The analysis was also limited to Einstein backgrounds, which nearly solve the equations of motion, but also in this way is more general than the analysis in I, which was limited to maximally symmetric backgrounds.

We have obtained formulas for the one-loop divergences up to quadratic terms in the curvature. These are all the divergences that arise at one loop in $d = 4$. The method we have used is a one-loop approximation of the single-field approximation of the Exact RG equation for gravity, as first derived in [23] for the EH-case and then extended to HDG in [27, 28, 19], to 3- d topologically massive gravity in [31], and beyond the one-loop approximation in [32, 33, 34].

The coefficients are related to the beta functions for the couplings. Comparing the action (2.1) with our results (4.3), we find the beta functions for the dimensionless couplings (4.6) as

$$\begin{aligned}\beta_{\tilde{G}} &= (d-2)\tilde{G} + B_1\tilde{G}^2, \\ \beta_{\tilde{\Lambda}} &= -2\tilde{\Lambda} + B_1\tilde{\Lambda}\tilde{G} + \frac{A_1}{2}\tilde{G}, \\ \beta_{\tilde{\alpha}+\frac{1}{d}\tilde{\beta}} &= -(d-4)\left(\tilde{\alpha} + \frac{1}{d}\tilde{\beta}\right) + C_1.\end{aligned}\tag{7.1}$$

As already remarked, working on an Einstein space prevents us from disentangling the beta functions of $\tilde{\alpha}$ and $\tilde{\beta}$. On the other hand, even though we did not write a term $\gamma R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ in the action, the divergence D_1 gives rise to a beta function

$$\beta_{\tilde{\gamma}} = -(d-4)\tilde{\gamma} + D_1.\tag{7.2}$$

The signs of the coefficients imply that α , β , γ are asymptotically free in $d = 4$ whereas $\tilde{G}$ goes in the UV to a fixed point $\tilde{G} = -(d-2)/B_1$. In order for this nontrivial fixed point to make sense, B_1 must be negative. In spite of the non-universality of the beta functions, many calculations indicate that B_1 is indeed negative for the linear split. For example, B_1 in (4.13) is negative for $m = \omega = 0$ and a wide range of a and b . With the exponential parametrization discussed in subsection 4.6, B_1 in the EH limit given in (4.21) is negative for $3 \leq d \leq 13$. We refer the reader to the literature for a more detailed discussion.

The explicit coefficients of the divergences, in arbitrary dimension, gauge and parametrization, are too complicated to write, and we have exhibited only some special cases. The universal values agree with the literature. The divergences also agree, in $d = 4$ and in the EH limit, with earlier calculations in general gauges [36, 37].

Most of the general features noted in I persist in the theories considered here, as we have observed. In particular, one striking feature that we have checked in full generality is the

existence of a “duality” symmetry under the change of parametrization (1.6), or

$$\omega \rightarrow 1 - \omega ; \quad m \rightarrow -m - \frac{2}{d} .$$

As observed in I, this transformation preserves the dimension of the quantum field and is also an invariance of the functional measure.³

Whether this duality extends also to other classes of actions, to other backgrounds and to higher loops are all questions that we leave for further investigation. Also left for future work is the calculation of divergences in the unimodular case $m = -1/d$, which contains in particular the unique self-dual theory $\omega = 1/2$, $m = -1/d$.

Acknowledgment

This work was supported in part by the Grant-in-Aid for Scientific Research Fund of the JSPS (C) No. 16K05331. ADP is grateful to CNPq for financial support.

A Heat kernel coefficients for Lichnerowicz Laplacians

The heat kernel coefficients for the Lichnerowicz Laplacians acting on spin-0, spin-1 and spin-2 fields on an Einstein manifold are,

$$\begin{aligned} b_0(\Delta_L^{(0)}) &= 1, \\ b_2(\Delta_L^{(0)}) &= \frac{1}{6}R, \\ b_4(\Delta_L^{(0)}) &= \frac{1}{180}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + \frac{5d-2}{360d}R^2, \end{aligned} \tag{A.1}$$

$$\begin{aligned} b_0(\Delta_L^{(1)}) &= d-1, \\ b_2(\Delta_L^{(1)}) &= \frac{d-7}{6}R, \\ b_4(\Delta_L^{(1)}) &= \frac{d-16}{180}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + \frac{5d^2-67d+182}{360d}R^2, \\ b_0(\Delta_L^{(2)}) &= \frac{(d+1)(d-2)}{2}, \\ b_2(\Delta_L^{(2)}) &= \frac{d^2-13d-14}{12}R, \\ b_4(\Delta_L^{(2)}) &= \frac{d^2-31d+508}{360}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + \frac{5d^3-127d^2+592d+1804}{720d}R^2. \end{aligned} \tag{A.2}$$

These formulae can be obtained by the methods described for example in Appendix B of [25]. Here we do *not* take into account isolated modes that do not contribute to the fluctuation $h_{\mu\nu}$. Thus these formulae hold in the case when the manifold has no Killing or conformal Killing vectors, or else if the manifold is noncompact and has a continuous spectrum, so that the spurious isolated modes are of measure zero.

In order to get the heat kernel coefficients of shifted Lichnerowicz Laplacians $\Delta + aR$, one can use

$$b_0(\Delta + aR) = b_0(\Delta),$$

³We recall that the pairs of measures proposed in [35] are dual in this sense.

$$\begin{aligned}
b_2(\Delta + aR) &= b_2(\Delta) - aRb_0(\Delta), \\
b_4(\Delta + aR) &= b_4(\Delta) - aRb_2(\Delta) + \frac{1}{2}a^2R^2b_0(\Delta).
\end{aligned}
\tag{A.3}$$

References

- [1] N. Ohta, R. Percacci and A. D. Pereira, “Gauges and functional measures in quantum gravity I: Einstein theory,” JHEP **1606** (2016) 115 [arXiv:1605.00454 [hep-th]].
- [2] L. Modesto, “Super-renormalizable Quantum Gravity,” Phys. Rev. D **86** (2012) 044005 [arXiv:1107.2403 [hep-th]].
- [3] T. Biswas, E. Gerwick, T. Koivisto and A. Mazumdar, “Towards singularity and ghost free theories of gravity,” Phys. Rev. Lett. **108** (2012) 031101 [arXiv:1110.5249 [gr-qc]].
- [4] L. Modesto and L. Rachwal, “Super-renormalizable and finite gravitational theories,” Nucl. Phys. B **889** (2014) 228 [arXiv:1407.8036 [hep-th]].
- [5] K. S. Stelle, “Renormalization of Higher Derivative Quantum Gravity,” Phys. Rev. D **16** (1977) 953.
- [6] J. Julve, M. Tonin, “Quantum Gravity with Higher Derivative Terms,” Nuovo Cim. B **46** (1978) 137.
- [7] E.S. Fradkin, A.A. Tseytlin, “Renormalizable Asymptotically Free Quantum Theory Of Gravity,” Phys. Lett. B **104** (1981) 377; “Renormalizable asymptotically free quantum theory of gravity,” Nucl. Phys. B **201** (1982) 469.
- [8] I. G. Avramidi and A. O. Barvinsky, “Asymptotic Freedom In Higher Derivative Quantum Gravity,” Phys. Lett. B **159** (1985) 269.
I. G. Avramidi, “Covariant methods for the calculation of the effective action in quantum field theory and investigation of higher derivative quantum gravity,” hep-th/9510140.
- [9] A. Salam and J. A. Strathdee, “Remarks On High-Energy Stability And Renormalizability Of Gravity Theory,” Phys. Rev. D **18** (1978) 4480.
- [10] R. Floreanini and R. Percacci, “The Renormalization group flow of the Dilaton potential,” Phys. Rev. D **52** (1995) 896 [hep-th/9412181].
- [11] A. Bonanno and M. Reuter, “Modulated Ground State of Gravity Theories with Stabilized Conformal Factor,” Phys. Rev. D **87** (2013) 084019 [arXiv:1302.2928 [hep-th]].
- [12] I. L. Shapiro, “Counting ghosts in the ghost-free non-local gravity,” Phys. Lett. B **744**, 67 (2015) [arXiv:1502.00106 [hep-th]].
- [13] E. Tomboulis, “1/N Expansion And Renormalization In Quantum Gravity,” Phys. Lett. B **70** (1977) 361.
E. Tomboulis, “Renormalizability And Asymptotic Freedom In Quantum Gravity,” Phys. Lett. B **97** (1980) 77.
E. T. Tomboulis, “Unitarity in Higher Derivative Quantum Gravity,” Phys. Rev. Lett. **52** (1984) 1173.

- [14] P. D. Mannheim, “Solution to the ghost problem in fourth order derivative theories,” *Found. Phys.* **37** (2007) 532 [hep-th/0608154].
- [15] S. Mukohyama and J. P. Uzan, “From configuration to dynamics: Emergence of Lorentz signature in classical field theory,” *Phys. Rev. D* **87** (2013) 065020 [arXiv:1301.1361 [hep-th]].
- [16] A. Salvio and A. Strumia, “Agravity,” *JHEP* **1406** (2014) 080 [arXiv:1403.4226 [hep-ph]].
- [17] M. B. Einhorn and D. R. T. Jones, “Naturalness and Dimensional Transmutation in Classically Scale-Invariant Gravity,” *JHEP* **1503** (2015) 047, [arXiv:1410.8513 [hep-th]].
- [18] L. Alvarez-Gaume, A. Kehagias, C. Kounnas, D. Lüst and A. Riotto, “Aspects of Quadratic Gravity,” *Fortsch. Phys.* **64** (2016) 176 [arXiv:1505.07657 [hep-th]].
- [19] N. Ohta and R. Percacci, “Higher Derivative Gravity and Asymptotic Safety in Diverse Dimensions,” *Class. Quant. Grav.* **31** (2014) 015024 [arXiv:1308.3398 [hep-th]].
- [20] R. Percacci, M. J. Perry, C. N. Pope and E. Sezgin, “Beta Functions of Topologically Massive Supergravity,” *JHEP* **1403** (2014) 083 [arXiv:1302.0868 [hep-th]].
- [21] R. Percacci and G. P. Vacca, “Search of scaling solutions in scalar-tensor gravity,” arXiv:1501.00888 [hep-th].
- [22] H. Hata, T. Kugo and N. Ohta, “Skew Symmetric Tensor Gauge Field Theory Dynamically Realized in QCD U(1) Channel,” *Nucl. Phys. B* **178** (1981) 527.
T. Kugo and S. Uehara, “General Procedure of Gauge Fixing Based on BRS Invariance Principle,” *Nucl. Phys. B* **197** (1982) 378.
- [23] M. Reuter, “Nonperturbative evolution equation for quantum gravity,” *Phys. Rev. D* **57** (1998) 971 [hep-th/9605030].
- [24] D. Dou and R. Percacci, “The running gravitational couplings,” *Class. Quant. Grav.* **15** (1998) 3449 [hep-th/9707239].
- [25] A. Codello, R. Percacci and C. Rahmede, “Investigating the Ultraviolet Properties of Gravity with a Wilsonian Renormalization Group Equation,” *Annals Phys.* **324** (2009) 414 [arXiv:0805.2909 [hep-th]].
- [26] G. de Berredo-Peixoto and I. L. Shapiro, “Higher derivative quantum gravity with Gauss-Bonnet term,” *Phys. Rev. D* **71** (2005) 064005 [hep-th/0412249].
- [27] A. Codello and R. Percacci, “Fixed points of higher derivative gravity,” *Phys. Rev. Lett.* **97** (2006) 221301 [hep-th/0607128].
- [28] M. Niedermaier, “Gravitational Fixed Points from Perturbation Theory,” *Phys. Rev. Lett.* **103** (2009) 101303; “Gravitational fixed points and asymptotic safety from perturbation theory,” *Nucl. Phys. B* **833** (2010) 226.
- [29] G. de Berredo-Peixoto and I. L. Shapiro, “Conformal quantum gravity with the Gauss-Bonnet term,” *Phys. Rev. D* **70** (2004) 044024 [hep-th/0307030].
- [30] Y. Pang, “One-Loop Divergences in 6D Conformal Gravity,” *Phys. Rev. D* **86** (2012) 084039 [arXiv:1208.0877 [hep-th]].

- [31] R. Percacci and E. Sezgin, “One Loop Beta Functions in Topologically Massive Gravity,” *Class. Quant. Grav.* **27** (2010) 155009 [arXiv:1002.2640 [hep-th]]
- [32] O. Lauscher and M. Reuter, “Flow equation of quantum Einstein gravity in a higher derivative truncation,” *Phys. Rev. D* **66** (2002) 025026 [hep-th/0205062].
- [33] D. Benedetti, P. F. Machado and F. Saueressig, “Asymptotic safety in higher-derivative gravity,” *Mod. Phys. Lett. A* **24** (2009) 2233 [arXiv:0901.2984 [hep-th]]; “Taming perturbative divergences in asymptotically safe gravity,” *Nucl. Phys. B* **824** (2010) 168 [arXiv:0902.4630 [hep-th]].
- [34] K. Groh, S. Rechenberger, F. Saueressig and O. Zanusso, “Higher Derivative Gravity from the Universal Renormalization Group Machine,” *PoS EPS -HEP2011* (2011) 124 [arXiv:1111.1743 [hep-th]].
- [35] K. Fujikawa, “Path Integral Measure for Gravitational Interactions,” *Nucl. Phys. B* **226** (1983) 437.
K. Fujikawa and O. Yasuda, “Path Integral for Gravity and Supergravity,” *Nucl. Phys. B* **245** (1984) 436.
- [36] R. E. Kallosh, O. V. Tarasov and I. V. Tyutin, “One Loop Finiteness Of Quantum Gravity Off Mass Shell,” *Nucl. Phys. B* **137** (1978) 145.
- [37] M. Y. Kalmykov, “Gauge and parametrization dependencies of the one loop counterterms in the Einstein gravity,” *Class. Quant. Grav.* **12** (1995) 1401 [hep-th/9502152].
M. Y. Kalmykov, K. A. Kazakov, P. I. Pronin and K. V. Stepanyantz, “Detailed analysis of the dependence of the one loop counterterms on the gauge and parametrization in the Einstein gravity with the cosmological constant,” *Class. Quant. Grav.* **15** (1998) 3777 [hep-th/9809169].